arXiv:1704.06554v1 [math.NT] 11 Apr 2017

EXTENDIBILITY OF SOME P_k SETS

BILGE PEKER AND SELIN (INAG) CENBERCI

Abstract. A set of m distinct positive integers $\{a_1, ...a_m\}$ is called a Diophantine m-tuple if $a_i a_j + n$ is a square for each $1 \leq i < j \leq m$. The aim of this study is to show that some P_k sets can not be extendible to a Diophantine quadruple when $k = 2$ and $k = -3$ and also to give some properties about P_k sets.

INTRODUCTION

Let n be an integer and A be a set $\{a_1, a_2, a_3, ..., a_m\}$ with m different positive integers. This set has the property $D(n)$ if $a_i a_j + n$ is a perfect square for all $1 \leq i < j \leq m$. Such a set is called as a Diophantine m-tuple (with the property $D(n)$) or a P_n set of size m.

The problem extending of P_k sets is an old one dating from the times of Diophantus [3] . So, this problem has been studied extensively. Diophantus studied finding four numbers such that the product of any two of them, increased by 1, is a perfect square. In [2, 4, 9], the more general form of this problem was considered.The most famous result in this area is the one found by Baker and Davenport [1] who proved for the four numbers $1, 3, 8, 120$ the property that the product of any two, increased by 1, is a perfect square. Dujella [4] has a vast amount of literature about this interesting problem.

Filipin [7] studied on the set $P_{-1} = \{1, 10, c\}$ and proved that there do not exist different positive integers $c, d > 1$ such that the product of any two distinct elements of the set $\{1, 10, c, d\}$ diminished by 1 is a perfect square.

In 1985, Brown [2], Gupta and Singh [8], Mohanty and Ramasamy [11] proved independently that if $n \equiv 2 \pmod 4$ then there does not exist a P_n set of size 4. Peker et al. [16] showed that for an integer k such that $|k| \geq 2$, the $D(-2k+1)-$triple $\left\{1, k^2, k^2 + 2k - 1\right\}$ can not be extended to a $D(-2k+1)-$quadruple.

Mohanty and Ramasay [10] showed non-extendibility of the set $P_{-1} = \{1, 5, 10\}$. Kedlaya [9] gave a list of non-extendibity of P_{-1} triples $\{1, 2, 45\}$, $\{1, 2, 4901\}$, $\{1, 5, 65\}$, $\{1, 5, 20737\}$, $\{1, 10, 17\}$ and $\{1, 26, 37\}$. Both Thamotherampillai [17] and Mootha & Berzsenyi [13] proved the non-extendibility of the set $P_2 = \{1, 2, 7\}$ by using different methods. Ozer [15] gave some P_{11} and P_{-11} triples.

Diophantine triples are classified as regular or irregular according to whether they provide the condition below.

Definition 1. [6] *A P_k-triple $\{a, b, c\}$ is called regular if it satisfies the condition $(c - b - a)^2 = 4(ab + k)$. This equation is symmetric under permutations of a, b, c.*

Date: April 4, 2017.
2000 *Mathematics Subject Classification.* Primary 11D09; Secondary 11D25.
Key words and phrases. Diophantine m-tuples, Pell Equation, P_k sets.

Definition 2. [12] *If p is an odd prime and $(a,p)=1$ then $\left(\frac{a}{p}\right)$ is defined as Legendre Symbol and* $\left(\frac{a}{p}\right)=\begin{cases} +1, & if \quad aRp \\ -1, & if \quad aNp \end{cases}$

Theorem 1. [12] *(Euleur Criterion) If p is a prime and p does not divide a positive integer a, then $\left(\frac{a}{p}\right) \equiv a^{\frac{p-1}{2}} \pmod p$.*

In this paper, we give some non-extendable P_2 and P_{-3} sets by using solutions of simultaneous Pell equations.

Theorem 2. *The set $P_2=\{7,14,41\}$ is non-extendable.*

Proof. Assume that the Diophantine triple $P_2=\{7,14,41\}$ with the property $D(2)$ can be extended with m to a Diophantine quadruple. *Then there exist integers x,y,z such that*

$$7m+2=x^2 \qquad (1)$$

$$14m+2=y^2 \qquad (2)$$

$$41m+2=z^2 \qquad (3)\,.$$

By eliminating m from the equations (1) and (2), we get

$$y^2-2x^2=-2$$

and this gives

$$2(x^2-1)=y^2 \qquad (4)\,.$$

In the last equation, since the left hand side is even, the right hand side must be even, too. So, we can write $y=2y_1$ $(y_1\in\mathbb{Z})$. Then the last equation becomes

$$x^2-1=2y_1^2.$$

From this, we can conclude that x must be odd, that is, $x=2x_1+1$ $(x_1\in\mathbb{Z})$. Then, the last equation will be as follows

$$2x_1\,(x_1+1)=y_1^2.$$

Since *the left hand side of this equation has multiple of 2, the right hand side must be even. So, we can write $y_1=2y_2$ $(y_2\in\mathbb{Z})$.* Now, we have $y=4y_2$. *If we write $y=4y_2$ in the equation (4), we find*

$$x^2-8y_2^2=1.$$

The fundamental solution of this Pell equation is $(x,y_2)=(3,1)\in\mathbb{Z}^2$. So, all solutions of this Pell equation can be given in the form $x_n+\sqrt{8}(y_2)_n=(3+\sqrt{8})^n$ by the usual methods. We get the general recurrence relation for the solutions of x as $x_n=6x_{n-1}-x_{n-2}$. Using this general recurrence relation, we can get some values of m from the equation (1). For these values of m, one can easily check that none of these values give any square of an integer for the equation (3). So, we can not get an integer solution for z. This means that the set $P_2=\{7,14,41\}$ is non-extendable. □

Remark 1. *A P_2-triple $\{7,14,41\}$ is called regular.*

Now, in the following theorem we show that another P_2 triple, namely $\{1, 7, 14\}$ can not be extended to a Diophantine quadruple by using above method.

Theorem 3. *The set $P_2 = \{1, 7, 14\}$ is non-extendable.*

Proof. Assume that there is a positive integer m and the set $P_2 = \{1, 7, 14\}$ can be extended with m. Then, let us find an integer m such that,

$$m + 2 = x^2 \qquad (5)$$

$$7m + 2 = y^2 \qquad (6)$$

$$14m + 2 = z^2 \qquad (7)\,.$$

where x, y, z are some integers satisfying the above equations.

By eliminating m from the equations (6) *and* (7) *we get*

$$z^2 - 2y^2 = -2$$

and this yields

$$2(y^2 - 1) = z^2 \qquad (8)\,.$$

In the last equation, since the left side is even, the right side must be even, too. So, we can write $z = 2z_1$ $(z_1 \in \mathbb{Z})$, then the equation (8) *becomes*

$$y^2 - 1 = 2z_1^2.$$

From this, we can conclude that y must be odd, that is, $y = 2y_1 + 1$ $(y_1 \in \mathbb{Z})$. Then, we find

$$2y_1\,(y_1 + 1) = z_1^2.$$

Since the *left hand side of this equation has multiple of* 2, *the right hand side must be even. So, we can write $z_1 = 2z_2$ $(z_2 \in \mathbb{Z})$.*

Consequently, $z = 4z_2$. If we write $z = 4z_2$ in the equation (8), *we find*

$$y^2 - 8z_2^2 = 1.$$

The fundamental solution of this Pell equation is $(y, z_2) = (3, 1) \in \mathbb{Z}^2$. So, all solutions of this Pell equation can be given in the form $y_n + \sqrt{8}(z_2)_n = (3 + \sqrt{8})^n$ by the usual methods. We get the general recurrence relation for the solutions of y as $y_n = 6y_{n-1} - y_{n-2}$. Using this general recurrence relation, we can get some values of m from the equation (6). *For these values of m, one can easily check that none of these values give any square of an integer for the equation* (5). *So, we can not get an integer solution for x. This means that the set $P_2 = \{1, 7, 14\}$ is non-extendable.* □

Remark 2. *A P_2-triple $\{1, 7, 14\}$ is called regular.*

In addition, we can give another P_2 triple which can not be extended to a quadruple.

Theorem 4. *The set $P_2 = \{41, 239, 478\}$ is non-extendable.*

Proof. Assume that there is a positive integer m *and the set* $P_2 = \{41, 239, 478\}$ *can be extended with* m*. Then, let us find an integer* m such that,

$$41m + 2 = x^2 \qquad (9)$$

$$239m + 2 = y^2 \qquad (10)$$

$$478m + 2 = z^2 \qquad (11)\,.$$

where x, y, z are some integers satisfying the above equations.

By eliminating m *from the equations* (10) *and* (11)*, we get*

$$2y^2 - z^2 = 2$$

and

$$2(y^2 - 1) = z^2.$$

Since this equation is the same, equation (4) *and* (8)*, in the next steps, we will apply similar methods done in Theorem 1 and 2. Hence, this completes the proof.* □

Remark 3. *A* P_2*-triple* $\{41, 239, 478\}$ *is called regular.*

Remark 4. *There is no set* P_2 *including any positive multiple of* 3*.*

Let's assume that $3k$ $(k \in \mathbb{Z}^+)$ *is an element of the set* P_2*. For any* $t \epsilon P_2$*, the equation*

$$3kt + 2 = x^2$$

must be satisfied. In modulo3, the following equation is deduced,

$$x^2 \equiv 2 \pmod 3.$$

Since 3 *is an odd prime and* $(2, 3) = 1$*, using Legendre Symbol and Euleur Criterion, we get* $\left(\frac{2}{3}\right) = -1$ *which means that the last equation is unsolvable. Therefore,* $3k$ $(k \in \mathbb{Z}^+)$ *can not be an element of the set* P_2*.*

Theorem 5. *The* P_{-3} *set* $\{3, 4, 13\}$ *can not be extendible.*

Proof. Assume that there is a positive integer m and the set $P_{-3} = \{3, 4, 13\}$ can be extended with m. Then let us find integers x, y, z such that,

$$3m - 3 = x^2 \qquad (12)$$

$$4m - 3 = y^2 \qquad (13)$$

$$13m - 3 = z^2 \qquad (14)$$

where x, y, z are some integers satisfying the above equations. Eliminating of m between (12), (13) yields

$$3y^2 - 4x^2 = 3 \qquad (15)$$

and then we have to abtain following equation

$$3(y^2-1)=4x^2 \qquad (16)$$

If we consider equation (16), since left hand side is divided 3, right hand side must be divided 3 too. So $3 \mid x^2$ and consequently $3 \mid x$. And we write $x=3x_1$ $(x_1 \in \mathbb{Z})$. Byusing this relation if we rewrite equation (16) ,we get

$$y^2-1=12x_1^2 \qquad (17)$$

this equation gives us y must be odd. Therefore $y=2y_1+1$ $(y_1 \in \mathbb{Z})$ and from last equation we obtain the last equation,

$$y_1\left(y_1+1\right)=3x_1^2. \qquad (18)$$

Left hand side of the equation (18) is the multiplication of two consecutive numbers. And we know that multiplication of two consecutive numbers is always even. So the right hand side must be even, that is $x_1=2x_2$ $(x_2 \in \mathbb{Z})$. We obtain from this $x=6x_2$, and the equation (18) turn such Pell equation

$$y^2-48x_2^2=1. \qquad (19)$$

This Pell equations' main solution is $(x_2,y)=(1,7) \in \mathbb{Z}^2$. All of the solutions of this Pell equation can be find such that $y_n+\sqrt{48}\left(x_2\right)_n=\left(7+\sqrt{48}\right)^n$, $n=0,1,2,...$ .For y solutions, the general recurrence relation is $y_{n+2}=14y_{n+1}-y_n$. For considering equations $(12),(13),(14)$ we can not find a m value, this gives our assumption is not true. So the P_{-3} set $\{3,4,13\}$ can not be extended. This completes the proof. □

Theorem 6. *The set $P_2=\{7,41,82\}$ can not be extendible.*

Proof. Assume that there is a positive integer m and the set $P_2=\{7,41,82\}$ can be extended with m. Then let us find integers x,y,z such that

$$7m+2=x^2 \qquad (20)$$

$$41m+2=y^2 \qquad (21)$$

$$82m+2=z^2 \qquad (22).$$

By eliminating m from the equations (21) and (22) we get

$$2y^2-z^2=2$$

and

$$2(y^2-1)=z^2.$$

Since this equation is the same equation with (4) and (8), the next steps are same in Theorem 1 and 2 . Therefore this completes the proof. □

Theorem 7. *The set $P_2=\{41,82,239\}$ can not be extendible.*

Proof. Assume that there is a positive integer m an the set $P_2 = \{41, 82, 239, m\}$ *can be extendible with* m. *Then let us find an integer* m *such that*

$$41m + 2 = x^2 \qquad (23)$$

$$82m + 2 = y^2 \qquad (24)$$

$$239m + 2 = z^2 \qquad (25)\,.$$

where x, y, z *are some integers satisfying above equations. Eliminating* m *from the equations* (23) *and* (24) *yieldst*

$$y^2 - 2x^2 = -2$$

and then we obtain following equation

$$2(x^2 - 1) = y^2 \qquad (26)\,.$$

In the last equation, since the left side is even, the right side must be even too. So we can write $y = 2y_1$ $(y_1 \in \mathbb{Z})$, *last equation gives us*

$$x^2 - 1 = 2y_1^2.$$

From this, we can conclude that x *must be odd. If we put* $x = 2x_1 + 1$ $(x_1 \in \mathbb{Z})$, *then we find*

$$2x_1\,(x_1 + 1) = y_1^2.$$

The left hand side of this equation has multiple of 2, *that is, the right hand side must be even. So we can put* $y_1 = 2y_2$ $(y_2 \in \mathbb{Z})$. *Thus we can write* $y = 4y_2$. *Putting* $y = 4y_2$ *in the equation* (26) *yields*

$$x^2 - 8y_2^2 = 1.$$

Main solution of this Pell equation is $(x, y_2) = (1, 7) \in \mathbb{Z}^2$. *All of the solutions of this Pell equation are of the form* $x_n + \sqrt{8}(y_2)_n = (3 + \sqrt{8})^n$, $n = 0, 1, 2, 3, \ldots$. *For* x *solutions general recurrence relation is* $x_n = 6x_{n-1} - x_{n-2}$. *From the equations* $(23), (24), (25)$ *we can not find a* m *value. Hence, our assumption is not true. This completes the proof.* □

Remark 5. *There is no set not only* P_2 *but also* P_{-3} *including any positive multiple of 5.*

Assume that $5k$ $(k \in \mathbb{Z}^+)$ is an element of both the sets P_2 and P_{-3}. For any $t_1 \in P_2$ and $t_2 \in P_{-3}$, we can write the following equations,

$$5kt_1 + 2 = x^2 \qquad \text{and} \qquad 5kt_2 - 3 = y^2.$$

From these equations, one can deduce in modulo 5 the followings

$$x^2 \equiv 2 \pmod 5 \qquad \text{and} \qquad y^2 \equiv 2 \pmod 5\,. \qquad (27)$$

Since 5 is an odd prime and $(2, 5) = 1$, using from Legendre Symbol, we get $\left(\frac{2}{5}\right) = -1$ which means that the equation (27) is unsolvable. Therefore, $5k$ $(k \in \mathbb{Z}^+)$ can not be an element of both the sets P_2 and P_{-3}.

References

[1] A. Baker, H. Davenport, *The equations* $3x^2-2=y^2$ *and* $8x^2-7=z^2$, Quarterly Journal of Mathematics, Oxford (2), 20 (1969), 129-137.

[2] E. Brown, *Sets in which* $xy+k$ *is always a square*, Math. Comp. 45 (1985), 613-620.

[3] L.E. Dickson, *History of theory of numbers,* Chelsea New York, 2, (1996).

[4] A. Dujella, *Diophantine m-tuples*, http://www.math.hr/∼duje/dtuples.html

[5] A. Dujella, *Diophantine quadruples and quintuples modulo 4*, Notes on Number Theory and Discrete Math. 4 (1998), 160-164.

[6] A. Dujella, A. Jurasic, *Some Diophantine triples and quadruples for quadratic polynomials*, J. Comb. Number Theory 3 (2) (2011), 123-141.

[7] A. Filipin, *Nonextendibility of* $D(-1)$*-triples of the form* $\{1,10,c\}$, Internat. Math. Math Sci. 35 (2005), 2217-2226.

[8] H. Gupta, K.Singh, *On k-triad sequences,* Internat. Journal Math. Math. Sci. 5 (1985), 799-804.

[9] K.S. Kedlaya, *Solving constrained Pell equations*, Mathematics of Computation 67, 222, (1998), 833-842.

[10] S. P. Mohanty, A. M. S. Ramasamy, *On* $P_{r,k}$ *sequences*, Fibonacci Quart. 23 (1985), 36-44.

[11] S. P. Mohanty, A. M. S. Ramasamy, *The simultaneous Pell equations* $5y^2-20=x^2$, $2y^2+1=z^2$, Journal Number Theory 18 (1984), 356-359.

[12] Mollin R.A. Fundamental Number Theory with Applications, CRC Press, 2008.

[13] V.K.Mootha and G.Berzsenyi, Characterizations and extendibility of P_t sets, Fibonacci Quartely, 27 (1989), 287-288.

[14] F.A.Muriefah, A. Rashed, *The simultaneous Diophantine equations* $y^2-5x^2=4$, $z^2-442x^2=441$,The Arabian Journal for Sci. 33 (2004), 1737-1746.

[15] O. Ozer, *A note on the particular set with size three*, Boundary Field Problems and Computer Simulation 55 (2016), 56-59.

[16] B. Peker, A. Dujella and S.I. Cenberci, *The non-extensibility of* $D(-2k+1)-$*triples* $\left\{1,k^2,k^2+2k-1\right\}$, Miskolc Mathematical Notes, 1, 16, (2015), 385-390.

[17] N. Thamotherampillai, *The set of numbers* $\{1,2,7\}$, Bull. Calcutta Math. Soc. 72 (1980), 195-197.

Mathematics Education Programme, Ahmet Keleşoğlu Education Faculty, Necmettin Erbakan University, Konya, Turkey.

E-mail address: bilge.peker@yahoo.com

Mathematics Education Programme, Ahmet Keleşoğlu Education Faculty, Necmettin Erbakan University, Konya, Turkey.

E-mail address: scenberci@konya.edu.tr